

INDICATIONS: For use in cattle as an aid in treating uncomplicated milk fever (parturient paresis).

DOSAGE AND ADMINISTRATION:

For intravenous administration only.
Cattle 250-500 mL
Dosage may be repeated at 8 to 12 hour intervals if required.

CAUTION: Solution should be warmed to body temperature and administered slowly. Entire contents should be used upon entering, as this product contains no preservative. Discard any unused portion. Do not use if solution is cloudy or contains precipitate.

NOTE: Monitor animal's condition closely. Re-treatment may be necessary. If there is no noticeable improvement within 24 hours following treatment, consult your veterinarian. Aseptic precautions should be observed.

Manufactured by
Nova-Tech Inc.
Grand Island, NE 68801

RMS 92-593
18-802-50

Iss. 10-19

Lot No.

Exp. Date

NDC 57319-630-07

Calcium Gluconate 23%

STERILE – PRESERVATIVE FREE

**FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN**

Net Contents: 500 mL



Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

EACH 500 mL CONTAINS:

Active Ingredients

Calcium Gluconate 23% w/v
(provides 10.71 g of Calcium* per 500 mL)
Water for Injection q.s.

Inactive Ingredients

Boric Acid
(as a solubilizing agent) 3.5 grams
Electrolytes per 1000 mL:
Calcium 1069 mEq;
Hypertonic pH 2.5-4.5

Store between 15°C–30°C (59°F–86°F).



TAKE TIME

OBSERVE LABEL
DIRECTIONS



Trademarks are property
of Clipper Distributing
Company, LLC