

**INDICATIONS:** For use as an aid in the treatment of actinomycosis (lumpy jaw), actinobacillosis (wooden tongue), and necrotic stomatitis in cattle.

**CONTRAINDICATIONS:** The use of sodium iodide is contraindicated in pregnancy and hyperthyroidism.

**DOSAGE AND ADMINISTRATION:** Using aseptic procedures, administer slowly by intravenous injection. Inject carefully to avoid deposition outside of the vein. The usual dose is 30 mg per pound of body weight (15 mL/100 lb). May be repeated at weekly intervals, if necessary.

**WARNING:** Not for use in lactating dairy cows.

**CAUTION:** Animals vary in their susceptibility of iodides. Administer with caution until the animal's tolerance is determined. Discontinue treatment and consult veterinarian if adverse reactions occur.

18-819-25  
RMS 92-1042

Iss. 04-12

Lot No.

Exp. Date

NDC 57319-601-06

# Sodium Iodide 20% Injection

**FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Net Contents: 250 mL**

 **PHOENIX™**

Manufactured for:  
Clipper Distributing Company, LLC  
St. Joseph, MO 64507

**COMPOSITION:** Each 100 mL of sterile aqueous solution contains:  
Sodium Iodide ..... 20 grams  
Water for injection ..... q.s.

**Store between 15°C and 30°C  
(59°F and 86°F).**

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Trademarks are property  
of Clipper Distributing  
Company, LLC

Manufactured by  
Nova-Tech Inc.  
Grand Island, NE 68801



3